

原文題目(出處)：	A rapidly growing gingival mass. Oral Surg Oral Med Oral Pathol Oral Radiol 2013;115:2-8
原文作者姓名：	HyukIl Kwon, Nam Hee Kim, PhD, Yonghoon Cha, Hyung Jun Kim, Ho-Gul Jung, Ja Seung Koo, Jong In Yook, Hyun Sil Kim
通訊作者學校：	Chonnam National University, Yonsei University Gwangju and Seoul, Korea
報告者姓名(組別)：	杜孟璇 (Intern I 組)
報告日期：	102.5.7

內文：

CLINICAL PRESENTATION

■ Present illness

A 71-year-old Korean woman was referred to the Department of Oral and Maxillofacial Surgery with a 10-day history of a rapidly growing mass occupying the oral cavity, which was associated with pain and spontaneous bleeding.

■ Clinical examination

- soft or slightly firm pedunculated mass
- approximately 4 X 5 X 5 cm in size
- on the anterior aspect of the lower gingiva, extended to the buccal and lingual vestibule, floor of mouth, and abutted the tongue
- not lobulated and was well vascularized, showed a reddish purple color, and had a smooth, shiny surface with no ulceration
- paresthesia of the lips or chin (-), palpable cervical lymph nodes (-)

■ Medical history

- controlled hypertension and diabetes mellitus.
- had been diagnosed with asthma
- had suffered a cerebrovascular accident but has completely recovered
- no history of alcohol intake or smoking

■ Laboratory tests and diagnostic

To trace the etiology of the rapidly growing gingival mass and to evaluate the operability, all the laboratory tests and diagnostic workup were performed after she was admitted.

- Routine complete blood count and blood chemistry tests: within normal limits
- Coagulation blood tests and urinalysis: no other noticeable findings
- Dental panoramic radiography: demonstrated slight bone destruction with an irregular border, which was confined to the lower anterior alveolar bone. There was neither diffuse basal bone destruction, nor periosteal reaction (Figure 1).




Fig. 1. Panoramic view demonstrates multiple root tips of the teeth and a destructive bone lesion (arrow) with an irregular border on the lower anterior alveolar bone. Periosteal reaction is not found.

To minimize the interval between biopsy and surgical operation, and the possibility of spreading cancer cells during the surgical intervention, the radiological evaluations were performed before the biopsy.

- Computed tomography (CT): the gingival mass extended anteriorly and displaced the buccinator muscle outward. The mass also extended posteriorly and approximated the tongue (Figure 2, A and B).




Fig. 2. Computed tomography imaging. A, Axial soft tissue window setting. At the level of roots of the lower anterior teeth, axial CT image demonstrates a soft tissue mass (asterisk) approximately 4 X 5 X 5 cm on the lower anterior gingiva. B, Axial bone window setting. The mass (asterisk) causes irregular alveolar bone destruction and extends anterioposteriorly.

- Magnetic resonance imaging (MRI) : revealed a mass approximately 5 cm in size on the anterior gingiva showing low signal intensity on T1- or T2-weighted images (Figure 3, A and B). On T1-weighted images with gadolinium (GD) enhancement (顯影劑增強), however, the mass showed high signal intensity with focal heterogeneity (Figure 3, C). The mass eroded alveolar bone and extended to the buccal and lingual vestibule, floor of mouth, and abutted the tongue on sagittal T1 with GD contrast images (Figure 3, D). Enlarged cervical lymph nodes were identified on neither CT nor MRI scans (Figures 2 and 3). Based on clinical examination and radiologic imaging, the initial impression was that of a primary

well-vascularized soft tissue tumor, such as a malignant vascular tumor, or a metastatic cancer.




Fig. 3. Magnetic resonance imaging. **A**, Axial T1-weighted image. **B**, Axial T2-weighted image. **C**, Axial T1 with gadolinium (GD) contrast. There is evidence of contrast enhancement of the mass, clearly defining a discrete mass (*asterisk*), which could not be seen distinctly on noncontrast-enhanced T1- or T2-weighted images. **D**, Sagittal T1 with GD contrast images demonstrate that the gingival mass slightly erodes alveolar bone only. The basal bone portion under the alveolus is intact. The tumor extends to the buccal and lingual vestibule, floor of mouth, and abuts the tongue (*asterisk*).

Initial biopsy was performed from the lower anterior gingiva by an oral and maxillofacial surgeon. The specimen was sent for histopathological evaluations to an oral pathologist.

DIFFERENTIAL DIAGNOSIS

The gingival tumor presented an admixture of small cells with scant cytoplasm and hyperchromatic round to oval nuclei, and large cells with abundant cytoplasm and peripheral large nuclei (Figure 4, A).




Fig. 4. Histopathologic findings. **A**, The gingival tumor shows an admixture of small cells and large cells with frequent mitoses. **B**, A tumor from the axilla shows nests of small cells (hematoxylin and eosin stain; original magnification $\times 200$).

The tumor was well vascularized, but there was no histologic evidence of malignant vascular tumor. Based on the light microscopic findings, differential diagnosis was needed for Ewing sarcoma (ES)/primitive neuroectodermal tumor (PNET), primary Merkel cell carcinoma (MCC), lymphoma, and combined small cell carcinoma (SCLC), which could be metastatic.

- Among them, ES/PNET was discounted, as it occurs most frequently in teenagers, and the probability of its occurring in our elderly patient was low. Besides, the present lesion showed no “onion skin”-type periosteal reaction in radiologic findings, often associated with ES/PNET.
- Primary MCC is a type of neuroendocrine tumor of the head and neck, as is SCLC. Primary MCC, lymphoma, and metastatic SCLC could not be discriminated only by microscopic findings, because they show similar histologic features. *Thyroid transcription factor 1 (TTF-1)* was used as a helpful marker to verify that the gingival lesion was the primary MCC or metastatic SCLC. Lymphoma could be ruled out simultaneously if the tumor cells could show the positivity for TTF-1.
- SCLCs are thought to originate from neuroendocrine cells in the bronchus. Hence, the diagnosis of SCLC is increasingly supported by a variety of neuroendocrine markers, such as chromogranin A, CD57, synaptophysin, neuron-specific enolase, and most recently CD56. Among the neuroendocrine markers, CD56 is widely used because of its high sensitivity compared with

other antibodies.¹ For the diagnosis of the present case, CD56 could be used as a significant tool in revealing the neuroendocrine origin of the gingival mass.

Metastatic tumor was suspected initially. The estimation was based on the following clinical evidence.

- First, the present gingival mass grew extremely rapidly, beyond comparison with the common malignant tumors of oral cavity origin.
- Second, the jawbone destruction by the tumor was confined to the alveolar bone, which implies that the oral cavity lesion was not a central lesion.

Still some conflict remained:

- the gingiva is not a frequent site of metastatic tumors
- Most metastatic tumors to the oral cavity are to the jawbones.
- The posterior areas of the mandible are the most frequent sites of cancer metastasis, because they contain hematopoietically active marrow; however, the abundant capillary network of chronically inflamed gingiva can entrap cancer cells, and can also become sites of metastatic cancer.²
- Metastatic tumors to the oral region account for approximately 1.0% to 1.5% of all malignant oral tumors.³ Moreover, metastatic tumors to the oral mucosa are more rarely reported than those in the jaw-bones by a ratio of 1.0:2.5.² The most common primary sites of metastatic oral cancers are the lungs, breasts, kidneys, bone, and colorectum, respectively. The breasts are the most common primary site for tumors that metastasize to the jawbones, whereas the lungs are the most common source for cancers that metastasize to the oral soft tissues.²

The patient merely displayed alveolar bone erosion without mandibular bone marrow infiltration by the tumor. Therefore, we could hypothesize that lung cancer had metastasized to the inflamed gingival tissue first and then infiltrated well-vascularized periodontal ligament spaces; however, our patient showed no symptoms associated with a lung lesion.

Lung cancers are classified according to histologic type.

This classification has important implications for clinical management and prognosis of the disease. The histologic subtypes of lung cancer are classified into 2 main categories:

- SCLC
- non-small cell lung carcinoma (NSCLC): include adenocarcinoma, squamous cell carcinoma, and large cell carcinoma.

NSCLCs are usually detected before metastasis or local spread occurs. On the other hand, SCLCs have invariably spread when they are first detected, even if the primary tumor appears small and localized. Our patient showed no previous pulmonary symptoms when the oral cavity tumor was first found, which may support the possibility of the metastatic SCLC.

Taking the clinico-histopathologic evaluation of the patient and epidemiologic references into consideration, the preliminary diagnoses for the initial gingival lesions were summarized as primary MCC, lymphoma, and probably metastatic SCLC.

ADDITIONAL INFORMATION AND DIAGNOSIS

To determine whether the oral lesion was primary or metastatic, immunohistochemical staining for the gingival specimen was performed and, simultaneously, positron emission tomography-computed tomography (PET-CT) images for the patient were taken.

- immunohistochemical staining: the anti-TTF-1 (Dako, M3575, Glostrup, Denmark) and anti-CD56 (Dako, M7304, Glostrup, Denmark) antibodies were used. TTF-1 is exclusively expressed in the epithelium of the lung and thyroid gland; therefore, it can be a highly selective marker for tumors arising in these organs. The gingival tumor cells exhibited focal positivity for TTF-1, confirming their pulmonary origin (Figure 5, A). Therefore, lymphoma could be ruled out.




Fig. 5. Immunohistochemical staining with TTF-1. Both the specimen from the gingiva (A) and axilla (B) show focal positivity within the nuclei of the tumor cells (original magnification $\times 200$).

The tumor cells also showed strong positivity for CD56, the marker for neuroendocrine differentiation (Figure 6, A).




Fig. 6. Immunohistochemical staining with CD56. Both the specimen from the gingiva (A) and axilla (B) show strong positivity in the membrane of the tumor cells (original magnification $\times 200$).

Both MCC and metastatic SCLC show the common neuroendocrine differentiation, which could not be discriminated by CD56 immunohisto-chemistry. The possibility of primary MCC could also be excluded by TTF-1 positivity of the tumor. Despite similar histopathologic features to SCLC, MCC presents the opposite reactivity for TTF-1 immunohisto-chemistry. Cheuk et al. reported that 82.7% of pulmonary SCLCs were positive for TTF-1, whereas 42% of extrapulmonary SCLCs were positive, but extrapulmonary neuroendocrine tumor of MCC showed negativity for TTF-1 without exception.^{3,4}

- PET-CT scanning: revealed not only the oral cavity mass but also an infiltrative mass in the left upper lobe of the lung with contiguous invasion of mediastinal pleura and several seeding nodules. Enlarged lymph nodes with 18-fluorodeoxyglucose uptake were also noted in the left hilar, pretracheal, and axillary fossa, but bilateral cervical lymph nodes were intact (Figure 7).



Fig. 7. PET-CT shows a mass in the lungs with several seeding nodules near the left mediastinum, as well as the mass in the oral cavity (arrowhead). Enlarged lymph nodes were also noted in hilar, pretracheal, and axillary fossa on the left (arrow), but bilateral cervical lymph nodes were intact.

Both the gingival and lung masses were interpreted as malignant tumors, because they infiltrated adjacent normal tissues. If one was a metastatic lesion from the other distant site, which was the primary? In which direction did the metastasis occur? Metastasis from the lung to the gingiva was more probable than vice versa in the present case, judging from clinical evidence of a lack of metastatic cervical lymph nodes.

- Additional biopsy was performed on the left axilla. Based on its anatomic location, the axillary mass was considered a regional lymph node around the lungs, and was radiologically diagnosed as metastatic lung cancer. Therefore, a more dangerous bronchoscopic lung biopsy, in comparison with an axillary lymph node biopsy, was deemed unnecessary. The specimen was obtained through a sonogram-guided gun biopsy by a radiologist and sent for histopathological evaluations to a pathologist. Histologically, the tumor from the axillary lesion presented typical features of small cell carcinoma. The tumor cells had scant cytoplasm and hyperchromatic round to oval nuclei (Figure 4, B).




Fig. 4. Histopathologic findings. A, The gingival tumor shows an admixture of small cells and large cells with frequent mitoses. B, A tumor from the axilla shows nests of small cells (hematoxylin and eosin stain; original magnification $\times 200$).

Immunohistochemical staining was performed for the axillary mass and compared with the results of the gingival mass. The same antibodies were used for the immunohistochemical staining. The tumor cells of both locations showed similar expression patterns of TTF-1 and CD56. They exhibited focal positivity for TTF-1 (Figure 5, A and B), and a strong positivity for CD56 (Figure 6, A and B).




Fig. 5. Immunohistochemical staining with TTF-1. Both the specimen from the gingiva (A) and axilla (B) show focal positivity within the nuclei of the tumor cells (original magnification $\times 200$).




Fig. 6. Immunohistochemical staining with CD56. Both the specimen from the gingiva (A) and axilla (B) show strong positivity in the membrane of the tumor cells (original magnification $\times 200$).

Results of immunohistochemical staining with TTF-1 offered the critical evidence to verify whether the gingival and lung masses were synchronous double primary tumors or the gingival and axillary masses were all metastatic lung cancer. Intriguingly, the gingival lesions showed different microscopic features from the axillary lesions. The gingival tumor presented an admixture of small cells and large cells (Figure 6, A), with peripheral large nuclei, prominent nucleoli, and abundant cytoplasm with well-defined cell borders. Large cells aggregated in a nodular pattern with interspersed components of small cells, and contained numerous clear cell-like portions.




Fig. 6. Immunohistochemical staining with CD56. Both the specimen from the gingiva (A) and axilla (B) show strong positivity in the membrane of the tumor cells (original magnification $\times 200$).

To determine the characterization for the cytoplasmic granules of the large cells observed in the gingival SCLC specimen, Periodic acid-Schiff (PAS) staining was performed. The large clear cells were positive for PAS staining (Figure 8, A) and were digested by diastase (澱粉糖化酵素) (Figure 8, B). These results revealed that the tumors showing clear cell changes contained abundant glycogen in the cytoplasm.




Fig. 8. PAS stain of gingival tumor. The tumor cells of the gingival SCLC shows positivity for PAS staining (A), which were digested by diastase (B) (original magnification $\times 200$).

Based on the clinico-pathologic analyses, the final pathologic diagnosis for both the axillary and gingival masses was metastatic SCLC. Histopathologically, the axillary mass was composed of small cell carcinoma, and the gingival mass was composed of combined small cell carcinoma.

After the diagnosis was made, the patient refused treatment and passed away about 2 months later.

DISCUSSION

- SCLC is a highly malignant neoplasm accounting for approximately 20% to 25% of all bronchial carcinomas and leads to a poor prognosis for patients, with high recurrence and a tendency to develop metastases widely throughout the body at an early stage in its clinical course.⁵⁻⁷
- According to the recent World Health Organization classification, when at least 10% of the tumor bulk of SCLC is made of non-small cell components—large cells, squamous cell carcinoma, or adenocarcinoma—it is ~~filed~~ as *combined-SCLC (c-SCLC)*.⁸
- c-SCLC is relatively rare, accounting for 2% to 14% of all cases of SCLC. Barnard⁹ first described SCLC as “oat cell carcinoma,” because of the ~~that~~ cell shape and scant cytoplasm. Barnard⁹ also described that there is cytologic heterogeneity within SCLC, describing “in all these tumors, cells other than small ‘oat’ cells could be found.” Abeloff et al.¹⁰ reported morphologic and biochemical changes of small cell carcinoma. According to this report, 5 of the 40 patients presented squamous cell carcinoma,¹⁰ adenocarcinoma,¹ and large cell carcinoma,¹ which developed concurrently with or after their diagnosis of SCLC.¹¹
- An in vitro study showed that SCLC containing a large cell component was less sensitive to radiation therapy and chemotherapy than pure SCLC, and several clinical studies suggested that patients with SCLC containing a large cell component showed a significantly poorer response to treatment and shorter survival time than those with pure SCLC.¹²⁻¹⁴ Thus, making an accurate pathologic distinction between pure SCLC and c-SCLC is important for therapeutic implications.
- In the present case, because the gingival mass was found before the lung mass, which accompanied no pulmonary symptoms, the diagnosis of metastatic SCLC of the gingiva was more of a challenge for pathologists. Worse still, the 2 biopsy specimens, from the gingival and axilla, showed different histomorphologic features, which made diagnosis even more difficult. To determine the reason for the difference in histomorphologic features between the 2 lesions, 2 hypotheses could be made:
 - the possibility of dual primary cancers
 - transformation during metastasisBecause the gingival tumor partly contained small cell portions, which were also observed in the axillary mass, the second hypothesis may be more reasonable than the former.
- Compared with the axillary tumor mass, the gingival tumor intriguingly contained large clear cells. From the result of PAS stain with digestion of diastase, it was determined that the clear cytoplasm contained carbohydrates,

such as glycogen, glycoprotein, and proteoglycans. This excluded adenocarcinoma, which contains mucin, from the differential diagnosis. Recent studies show that glycogen may be used by carcinoma cells as a source for energy because of the limited supply of nutrients and oxygen in a rapidly growing carcinoma.¹⁵ Metastases are complicated processes in which the tumor cells have to detach from the primary tumor, and then must spread in the tissue, invade the lymphovascular system, survive the journey in the circulation, and settle down at the metastatic site. The histomorphologically altered metastatic lesion associated with glycogen accumulation suggests that the tumor underwent the alteration of glucose metabolism during the strained metastatic process.

- In summary, metastatic SCLC with different subtypes is extremely rare, although metastases of small cell carcinoma occur frequently throughout the body. The present case shows different histopathological features in the metastatic lesions, which makes diagnosis more difficult. To reveal the primary site, careful clinico-pathologic examinations and proper immunohistochemical studies were required. Although there are no established informed data on the prognosis of c-SCLC associated with metastases, it would likely be poorer than that of common SCLC, according to our case.

題號	題目
1	Metastatic tumors to the oral region account for approximately _____ % of all malignant oral tumors. (A) 1% (B) 5% (C) 10% (D) 30%
答案(A)	出處：Oral and Maxillofacial Pathology, 3rd edition, p.563, Metastases to the oral soft tissue
題號	題目
2	The most common site for oral soft tissue metastases is _____. (A) Tongue (B) Mucosa (C) Gingiva (D) Lip
答案(C)	出處：Oral and Maxillofacial Pathology, 3rd edition, p.563, Metastases to the oral soft tissue